RESCEU-2/15

A new estimation method for mass of an isolated neutron star using gravitational waves

Kenji Ono,[1, 2, *] Kazunari Eda,[1, 3, †] and Yousuke Itoh[3, ‡]

[1] *Department of Physics, Graduate School of Science,*
The University of Tokyo, Tokyo, 113-0033, Japan
[2] *Institute for Cosmic Ray Research (ICRR), University of Tokyo,*
5-1-5 Kashiwanoha, Kashiwa, Chiba 277-8582, Japan
[3] *Research center for the early universe (RESCEU),*
Graduate School of Science, The University of Tokyo, Tokyo, 113-0033, Japan

We investigate a possibility of estimating mass of an isolated rapidly rotating neutron star (NS) from a continuous gravitational wave (GW) signal emitted by the NS. When the GW passes through the gravitational potential of the NS, the GW takes a slightly longer time to travel to an observer than it does in the absence of the NS. Such a time dilation effect holds also for photons and is often referred to as the gravitational time delay (or the Shapiro time delay). Correspondingly, the phase of the GW from the NS shifts due to the Coulomb type gravitational potential of the NS, and the resulting logarithmic phase shift depends on the mass, the spin frequency of the NS, and the distance to the NS. We show that the NS mass can, in principle, be obtained by making use of the phase shift difference between two modes of the continuous GW such as once and twice spin frequency modes induced by a freely precessing NS or a NS containing a pinned superfluid core. We estimate the measurement accuracy of the NS mass using Monte Carlo simulations and find that the mass of the NS with its ellipticity 10^{-6} at 1 kpc is typically measurable with an accuracy 20% using Einstein Telescope.

arXiv:1503.01231v1 [gr-qc] 4 Mar 2015

I. INTRODUCTION

Mass is a fundamental quantity of an astronomical object. In the case of a neutron star (NS), mass is important, among other aspects [1], as it gives us a clue to study the equation of state of dense matter beyond the nuclear density, which is still uncertain through theoretical studies or experiments on the Earth (See, e.g., [2]).

Almost all the measurements of NS masses so far are limited for NSs in binaries. An exception is for the nearby isolated NS RX J 185635-754 [3], where its radiation radius and gravitational redshift have been measured modulo detailed modeling of its atmosphere [4] (See also the comments mentioned in the Sec. 4 of [5] and Fig. 3 there). Since an isolated star may have a different evolutionary history from a star in a binary (namely due to their mutual mass transfer), the former may follow a different mass function than the latter and hence it is interesting to measure masses of as many isolated NSs as possible. This paper proposes a new method to measure a mass of an isolated rapidly rotating NS that emits gravitational wave (GW).

We expect direct detection of GWs within the next decade [6]. The second-generation gravitational wave detectors such as KAGRA [7], advanced Laser Interferometric Gravitationalwave Observatory (aLIGO) [8], and advanced Virgo (aVirgo) [9] are under construction. Conceptual studies of third-generation gravitational wave telescopes are also on-going [10, 11]. One of the most important family of GW sources for those GW detectors is a compact binary coalescence (CBC). Detection of GW from a CBC event enables us to measure masses of compact stars in the binaries to sub-percent level. A possibility is actively discussed where we can study internal structure of the compact stars (if they are regular ones) through the tidal disruption of the binary stars during the binary merger ([12] for a review). However, again, the estimates of masses possible from detection of GWs from CBC events are only for stars in binaries. We here focus on another type of GW sources, namely, isolated rapidly rotating non-axisymmetric NSs and show that we can estimate masses of isolated NSs by detection of those sources with one of proposed third-generation telescopes, namely, Einstein Telescope (ET) [10].

In fact, our method proposed in this paper utilizes the fact that phase of GW from an isolated rapidly rotating compact star is modulated by the Coulomb type gravitational field of the star itself. This logarithmic phase shift is well-known in quantum mechanics [13], and partly incorporated in the GW waveform from CBC [14]. This Coulomb type phase shift corresponds to the famous gravitational time delay (or the Shapiro time-delay when it is observed in a binary system). But this phase shift seems neglected so far in the context of the studies searching for GWs from isolated pulsars. Then our idea to extract mass $M_{\rm NS}$ of an isolated pulsar at distance r from GW observation is as follows. In the case of GW, the Coulomb phase shift take a form of

---

[*] Electronic address: kenji@icrr.u-tokyo.ac.jp
[†] Electronic address: eda@resceu.s.u-tokyo.ac.jp
[‡] Electronic address: yousuke˙itoh@resceu.s.u-tokyo.ac.jp

$\delta\Phi(\omega_{\rm gw}) = 2\omega_{\rm gw} M_{\rm NS} \ln(2\omega_{\rm gw} r)$ for GW with frequency $\omega_{\rm gw}$ [15]. If we detect two GW modes at frequencies $\omega_{\rm gw,1}$ and $\omega_{\rm gw,2}$ from the same pulsar, we could find the mass by $\delta\Phi(\omega_{\rm gw,1}) - K\delta\Phi(\omega_{\rm gw,2}) = 2\omega_{\rm gw,1} M_{\rm NS} \ln K$ where $K \equiv \omega_{\rm gw,1}/\omega_{\rm gw,2}$. Before examining the feasibility of this idea, we first mention whether a pulsar could emit continuous GW at (more than) two frequencies.

A rotating tri-axial object with its rotational frequency $\omega_{\rm rot}$ emits GW mainly at frequency $2\omega_{\rm rot}$ when the rotational axis of the object is aligned to the one of the principal axis of its moment of inertia. If the former axis is not aligned to any one of the latters and this object rotates as a rigid body as a whole, this object will freely precess and emits GW at frequencies close to $\omega_{\rm rot}$ and $2\omega_{\rm rot}$ [16]. When the former axis is not aligned to any one of the latters and this object has a pinned superfluid layer(s), Jones proposed a possibility that this object could steadily rotate at a constant frequency and emits GW at frequencies close to $\omega_{\rm rot}$ and $2\omega_{\rm rot}$ [17]. The difference between the two scenarios [16] and [17] is that the latter does not freely precess and hence it shows no electromagnetic signature of free precession.

Observational evidences of existence of freely precessing NSs are recently reported by Makishima *et al.* [18, 19] where they found phase modulations in pulsations of the magnetar 4U 0142+61 and 1E 1547.0–5408 using the Suzaku X-ray observatory. They interpreted that those phase modulations are due to free precessions of those objects. There are several other objects that show signatures of free precessions, but it is not clear that current theoretical understandings of NS interior structures allow a free precession sustainable long enough so that we have any practical chance to observe it at all (See, e.g., [20–22] for discussion and references therein). This in turn means that the findings by Makishima *et al.* and further detections of freely precessing NSs give us an insight onto the interior structure of NSs and drive our theoretical understandings on it.

Indeed, majority of pulsars do not show any clear signature of free precession. However, Jones pointed out that a NS may have a pinned superfluid. Jones then advocated a theoretical possibility that a NS could rotate steadily even when the axis of the moment of inertia of its solid outer crust is misaligned to its rotational axis [17]. As a result, such a NS emits GW at two frequencies close to $\omega_{\rm rot}$ and $2\omega_{\rm rot}$ without showing any electromagnetic signature of free precession. This proposal by Jones motivates Bejger and Królak [23] to study a search method for GWs from known pulsars at once and twice their spin frequencies. Motivated by these observations and the theoretical proposal by Jones, we here assume a pulsar that emits GW at frequencies $\omega_{\rm gw,1} \simeq \omega_{\rm rot}$ and $\omega_{\rm gw,2} = 2\omega_{\rm gw,1} \simeq 2\omega_{\rm rot}$ to study feasibility of our idea of estimating the pulsar mass using Coulomb phase shifts. The GW radiation at the frequency $\omega_{\rm gw,1} \simeq \omega_{\rm rot}$ is sometimes called a "wobble mode" or "wobble radiation" for a freely precessing NS. Here we would like to include a non-precessing NS that emits GW at that frequency, we here call this mode "the first harmonic mode" while the GW at 2ω "the second harmonic mode" (or "the first overtone mode"). Applications of our idea to GW modes other than the ones considered here or extensions to the case where multiple GW modes more than two are available should be straightforward.

The rest of this paper is organized as follows. In Sec. II, we construct a GW subject to a gravitational Coulomb phase shift due to the gravitational potential created by the GW source. In Sec.III, we apply our formulation to an isolated NS and examine measurement accuracies of the NS parameters. Summary and conclusion are presented in Sec.IV. Throughout this paper, we adopt the geometric unit system in which both the light speed c and the gravitational constant G are unity.

II. GW WAVEFORMS PROPAGATING THROUGH NS GRAVITATIONAL POTENTIAL

A. Green's function

In this section, we derive the explicit form of the gravitational phase shift for GW caused by the gravitational potential of the source itself. The derivation in this section is basically based on the section 2 in [15]. Consider an isolated GW source with mass M and located at $r = 0$. The dominant static part of the exterior gravitational field far away from this object is well approximated by

$$ds^2 = -\left(1 - \frac{2M}{r}\right) dt^2 + \left(1 + \frac{2M}{r}\right) \delta_{ij} dx^i dx^j . \tag{1}$$

A metric perturbation $\bar{h}^{\mu\nu} \equiv \eta^{\mu\nu} - \sqrt{-g} g^{\mu\nu}$ with respect to this metric satisfies the following wave equations

$$\left[-\left(1 + \frac{4M}{r}\right) \partial_t^2 + \boldsymbol{\nabla}^2\right] \bar{h}^{\mu\nu} = 0 \tag{2}$$

outside of the GW source in the harmonic gauge $\bar{h}^{\mu\nu}{}_{,\nu} = 0$ where $\eta^{\mu\nu}$ denotes the flat spacetime metric. Motivated by this observation and to capture the dominant effect of the gravitational field of the source onto the GW phase, we express the full Einstein equations including the source under the harmonic gauge condition in the following form,

$$\Box_{\rm M} \bar{h}^{\mu\nu} = -16\pi\tau^{\mu\nu} . \tag{3}$$

The effective energy-momentum pseudotensor $\tau^{\mu\nu}$ and the differential operator $\Box_{\rm M}$ are defined as

$$\Box_{\rm M} = \left[-\left(1+\frac{4M}{r}\right)\partial_t^2 + \boldsymbol{\nabla}^2 \right], \tag{4a}$$

$$\tau^{\mu\nu} = (-g)\left(T^{\mu\nu} + t^{\mu\nu}_{\rm LL} + t^{\mu\nu}_{\rm M}\right), \tag{4b}$$

$$t^{\mu\nu}_{\rm M} = -\frac{1}{16\pi}\left[\bar{h}^{\mu\alpha}{}_{,\beta}\bar{h}^{\nu\beta}{}_{,\alpha} + 2\bar{h}^{0j}\bar{h}^{\mu\nu}{}_{,0j} + \bar{h}^{jk}\bar{h}^{\mu\nu}{}_{,jk} + \left(\bar{h}^{00} - \frac{4M}{r}\right)\bar{h}^{\mu\nu}{}_{,00} \right], \tag{4c}$$

where $t^{\mu\nu}_{\rm LL}$ denotes the Landau-Lifshitz pseudotensor [24] and M denotes the total mass of the source. It should be noted that the term $(4M/r)\,\bar{h}^{\mu\nu}{}_{,00}$ appears in the both sides in Eq. (3). Because the 00 component of the metric perturbation is evaluated in the wave-zone as $\bar{h}^{00} = 4M/r$ to the lowest order in the weak field approximation, we intentionally added this term to the both sides in Eq. (3). Introducing the "potential barrier" $4M/r$ in the wave operator, we can take into account the deviation of the curved spacetime light cone from the flat spacetime one to the lowest order in the weak field approximation. As will be seen in the end of this section, this mass term brings the gravitational phase shift in the GW phase.

Equation. (3) can be formally solved by using the retarded Green's function $G^{(+)}_{\rm M}(x,x')$ as,

$$\bar{h}^{\mu\nu}(x) = -16\pi \int d^4x'\; G^{(+)}_{\rm M}(x,x')\,\tau^{\mu\nu}(x'), \tag{5}$$

where the retarded Green's function obeys the following equations.

$$\Box_{\rm M} G^{(+)}_{\rm M}(x,x') = \delta^{(4)}(x-x'), \tag{6a}$$

$$G^{(+)}_{\rm M}(x,x') = \sum_{\ell m}\int d\omega\; e^{i\sigma_\ell}{\rm sgn}(\omega)\left[\Psi^{+\omega\ell m}(x)\,\Psi^{S\omega\ell m*}(x')\,H(r-r') + \Psi^{S\omega\ell m}(x)\,\Psi^{+\omega\ell m*}(x')\,H(r'-r)\right]. \tag{6b}$$

The functions $\Psi^{+\omega\ell m}$ and $\Psi^{S\omega\ell m}$ are composed of the spherical Coulomb wave functions $u^{(+)}(\rho)$ and $F_\ell(\rho)$ [13]:

$$\Psi^{+\omega\ell m}(x) = \sqrt{\frac{|\omega|}{2\pi}}e^{-i\omega t}\rho^{-1}u^{(+)}_\ell(\rho)\,Y_{\ell m}(\theta,\phi), \tag{7a}$$

$$\Psi^{S\omega\ell m}(x) = \sqrt{\frac{|\omega|}{2\pi}}e^{-i\omega t}\rho^{-1}F_\ell(\rho)\,Y_{\ell m}(\theta,\phi), \tag{7b}$$

where $\sigma_\ell = \arg\Gamma(\ell+1-2iM\omega)$, $\rho = r\omega$, and $H(x)$ denotes the Heaviside step function. The radial functions $u^{(+)}_\ell(\rho)$ and $F_\ell(\rho)$ can be evaluated in the wave zone or the near zone as follows.

$$u^{(+)}_\ell(\rho) \to \exp\left[i\left(\rho + 2M\omega\ln 2\rho - \frac{1}{2}\ell\pi\right)\right] \qquad \text{for} \quad \rho\to\infty, \tag{8a}$$

$$F_\ell(\rho) \to c_\ell\rho^{\ell+1} \qquad \text{for} \quad \rho\to 0, \tag{8b}$$

with $c_\ell = 2^\ell e^{\pi M\omega}\left|\Gamma(\ell+1-2iM\omega)\right|/(2\ell+1)!$. Since the distance to the source is much greater than the size of the source $r \gg r'$, substituting Eqs.(7a)-(8b) into Eq.(6b) brings the Green's function to the following form,

$$G^{(+)}_{\rm M}(x,x') = \sum_\ell \frac{(-i)^\ell n^{\langle L\rangle}x'_{\langle L\rangle}}{2\,(2\pi\ell!)^2\, r}\int d\omega\; \Gamma(1+\ell-2iM\omega)\,e^{\pi M\omega}\omega^\ell e^{-i\left(t-r-2M\ln 2\omega r - t'\right)\omega} + \mathcal{O}\left(r^{-2}\right), \tag{9}$$

where the angle bracket attached to the indices denotes the symmetric-trace-free operation, the suffix L stands for ℓ tensorial products of vectors, and the unit vector n^j is defined as $n^j = x^j/r$. Combining Eqs. (5) and (9), we obtain the GW waveforms affected by the gravitational potential of the source. The authors of Ref. [15] then adopted slow motion approximation and expanded this Green's function in $M\omega$ and arrived at the GW waveform including the tail term which was first derived by Blanchet and Damour [14]. They then pointed out that the dominant part of the tail term originates from the deviation of the true light cone from the flat light cone due to the gravitational potential of the GW source itself, which appears as the phase shift $2iM\omega\ln 2\omega r + i\sigma_\ell$ in Eq. (9). Since we focus on this gravitational phase shift for GWs, we drop the $M\omega$ term in the amplitude but retain the $M\omega$ term in the phase up to the first order in our analysis.

Finally we note that this phase shift corresponds to the well-known gravitational time delay (or the Shapiro time delay in the case of a binary). The GW propagates along the true light cone of the curved spacetime, and so takes a slightly longer time to travel to the detector than it does in the absence of the source.

B. Continuous GWs through NS gravitational potential

Let us now assume an isolated rapidly rotating NS with mass $M_{\rm NS}$ as a GW source. As mentioned in the introduction, an isolated rapidly rotating NS would emit GWs at distinct frequencies at the same time. We define the volume integral of the stress-energy tensor T^{jk} as,

$$S^{jk}\left(t\right)=\int d^3x\ T^{jk}\left(t,\boldsymbol{x}\right). \tag{10}$$

If the source has different frequency modes $\omega_1,\omega_2,\cdots$, the Fourier component of $S^{jk}\left(t\right)$ can be decomposed into the sum of the delta function,

$$\tilde{S}^{jk}\left(\omega\right)=\sum_n 2\pi\tilde{S}^{jk}_n\delta\left(\omega-\omega_n\right), \tag{11}$$

where S^{jk}_n is the coefficient corresponding to the n-th mode. Combining Eqs. (5), (9)-(11), we arrive at the GW waveform including the gravitational phase shift in the quadrupole approximation,

$$\bar{h}^{jk}\left(t\right)=\frac{4}{r}\sum_n\tilde{S}^{jk}_n e^{-i\Psi_n(t)}, \tag{12a}$$

$$\Psi_n\left(t\right)=\left(t-r-2M_{\rm NS}\left[\ln(2r\omega_n)+C\right]\right)\omega_n. \tag{12b}$$

The constant C is the Euler's number.

It is easy to see that when $M_{\rm NS}\to 0$, Eq. (12a) with Eq. (12b) reduces to the well-known quadrupole gravitational waveform as expected. Inclusions of higher order mass multipoles or current multipoles, if necessary, are straightforward.

III. MEASUREMENT ACCURACY OF NS MASS

In this section, we apply the GW waveform including the gravitational phase shift effect to an isolated rapidly rotating NS that emits GWs at two frequencies, $\omega_{\rm gw,1}=\omega$ and $\omega_{\rm gw,2}=2\omega$.

A. GWs from an isolated rapidly rotating non-axisymmetric NS

A non-axisymmetric NS rotating around its principal axis produces the quadrupole GW radiation with a frequency $2\omega_{\rm rot}$, where $\omega_{\rm rot}$ denotes the rotational frequency of the NS. However, the angular momentum axis of the NS does not generally align with its principal axis because of, for example, the strong toroidal magnetic field inside the NS [25]. When the NS deviates from the spherical symmetry, the misalignment between the angular momentum axis and the principal axis in general leads to a free precession of the NS. It would also be possible that the NS steadily rotates at $\omega_{\rm rot}$ at the fixed rotational axis in an inertial frame if it has a pinned superfluid component inside it [17]. Those two kinds of NSs radiate the GW with two frequency modes $\omega_{\rm gw,1}=\omega$ and $\omega_{\rm gw,2}=2\omega$ which we call the first harmonic mode and the second harmonic mode, respectively [16, 17].

The GW waveforms in these two scenarios are generally different from each other for tri-axial NSs [16, 17, 23, 26, 27]. . Interestingly, in the case of bi-axial NSs, those two families of the GW waveforms take the same form (shown e.g., in [28]), which we adopt in this paper to apply our result to both scenarios.

We then adopt the following GW waveforms

$$h_+=A_{+,1}\cos\Psi_1+A_{+,2}\cos\Psi_2, \tag{13a}$$

$$h_\times=A_{\times,1}\sin\Psi_1+A_{\times,2}\sin\Psi_2 \tag{13b}$$

where A and Ψ denote the amplitude and the phase and are defined as

$$A_{+,1}=\frac{1}{4}h_0\sin 2\theta\sin\iota\cos\iota, \tag{14a}$$

$$A_{+,2}=\frac{1}{2}h_0\sin^2\theta\left(1+\cos^2\iota\right), \tag{14b}$$

$$A_{\times,1}=\frac{1}{4}h_0\sin 2\theta\sin\iota, \tag{14c}$$

$$A_{\times,2}=h_0\sin^2\theta\cos\iota, \tag{14d}$$

$$\Psi_1=\left(t-r-2M_{\rm NS}\ln 2r\omega\right)\omega+\varphi_0, \tag{14e}$$

$$\Psi_2 = 2\left(t - r - 2M_{\rm NS}\ln 4r\omega\right)\omega + 2\varphi_0. \tag{14f}$$

The angle ι is called the inclination defined as the angle between the rotational axis and the line-of-sight, and θ is the misalignment angle between the principal axis and the angular momentum axis. The angle φ_0 is the GW phase at the reference time $t = 0$ at the NS [33]. The overall amplitude h_0 is given by $h_0 = 4\varepsilon I\omega^2/r$ where I and ε are the principle moment of inertia and the ellipticity, respectively. Note that the log-term appears in the GW phase due to the gravitational potential of the NS.

B. How to extract the NS mass from GW signals

In this paper, we take an example of GWs from an isolated rapidly rotating non-axisymmetric NS to demonstrate the measurement accuracy of the NS mass. A NS would emit GWs of several modes such as the first and the second harmonic modes mentioned in this paper, wobble mode if the spin axis of the NS precesses, f-mode and r-mode if the corresponding stellar oscillations are excited. If the initial GW phases of two modes at the NS are known, or the ratio of the two GW phases is equal to the ratio of the GW frequencies of the two mode, the NS mass can be estimated by subtracting the GW phase of one mode from the other with the ratio multiplied appropriately. In the case of the first and second harmonic modes considered in this paper, this assumption is guaranteed because the both modes are generated by the deformation of the crust of the NS. Multiplying Ψ_1 by 2 Subtracting Eq. (14f) from Eq. (14e), we obtain

$$2\Psi_1 - \Psi_2 = 4M_{\rm NS}\omega\ln 2. \tag{15}$$

Equation. (15) indicates that the difference between the two modes separates the NS mass and the distance to the NS. Therefore, the NS mass alone can be estimated by GW observations. However, the accuracy of the constant phase is typically of the order of 0.1 for a signal to noise ratio (SNR) 10 even in targeted searches for known pulsars where $\omega_{\rm rot}(\simeq \omega_{\rm gw,1})$ is well-known by electromagnetic observations. Then, the second generation GW detectors would have a little chance of detecting the log-term appeared in Eq. (15) with enough sensitivity to estimate the NS mass. On the other hand, a third generation GW detector whose sensitivity is about an order of magnitude better than that of the second generation ones could determine the NS mass from the GW signal.

As will be discussed in the next section, we consider targeted searches for GWs from "known" rapidly rotating non-axisymmetric NSs by a third generation detector. Those objects are "known" in the sense that electromagnetic observations and/or the second generation detectors will have detected them before the search using third generation detectors. We then assume that the positions of the NSs in the sky and its GW frequencies (or rotation frequncies) and their times derivatives are known in advance. Those parameters are not search parameters in the following hypothetical search with a third generation GW detector.

C. Monte Carlo simulations

In this section, we investigate how accurately the mass of the NS can be determined by GW observations. Measurement errors are estimated by using the Fisher analysis which is briefly summarized in Appendix B. A GW signal from an isolated rapidly rotating non-axisymmetric NS is given by Eqs. (A1)-(A5). The GW signal is characterized by 6 waveform parameters $\boldsymbol{\lambda} = \{h_0, \theta, \iota, \psi, \phi_0, M_{\rm NS}\}$ in targeted searches. To demonstrate the accuracies of the waveform parameters $\boldsymbol{\lambda}$, we perform Monte Carlo simulations for the observation time $T_{\rm obs} = 3$ years in which we treat $\alpha, \delta, \theta, \iota$, and ψ as random variables. Here, the sky position of the NS is specified by two parameters, the right ascension α and the declination δ. The NS mass $M_{\rm NS}$ is assumed to be 1.4 $M_\odot$. We adopted $\varepsilon = 10^{-6}$ [29], $I = 10^{38}$ kg $\cdot$ m^2, and selected distances and frequencies to compute the amplitude h_0. For each simulation, we generate 10000 sets of random variables each of which is distributed according to the uniform distribution.

Figure. 1 displays the cumulative distribution functions of relative errors in the NS mass estimation $\Delta M_{\rm NS}/M_{\rm NS}$ from our Monte Carlo simulations for ET observations. We investigated the cumulative distribution functions for two different GW frequencies 300 Hz and 500 Hz. These figures show, for example, in the case of the GW signal with $f_{\rm rot} \simeq f_{\rm gw,1} = \omega_{\rm gw,1}/(2\pi) = 500$ Hz from NSs at $r = 1$ kpc in ET observations, that masses of about a 70 percent of isolated rapidly rotating non-axisymmetric NSs are measurable with an accuracy of $\Delta M_{\rm NS}/M_{\rm NS} \simeq 0.2$. It can be detected with SNR $= \mathcal{O}(10)$ by a single second generation detector because the sensitivity of ET is about an order of magnitude better than that of second generation one. Therefore, if an isolated rapidly rotating non-axisymmetric NS is detected by a network of second generation detectors, the mass of the NS can be determined with an accuracy of $\Delta M_{\rm NS}/M_{\rm NS} = \mathcal{O}\left(0.1\right)$ in a single third generation detector. Also, Fig.1 indicates that the accuracy of the NS mass becomes better and better as the GW frequency increases. This behavior can be traced to Eq. (15). The effect of the gravitational phase shift is proportional to the rotational frequency. So the NS mass can be more easily extracted from the GW signal with the higher frequency.

Figure. 2 shows that the scatter-plot of the NS mass accuracy $\Delta M_{\rm NS}/M_{\rm NS}$ as a function of the inclination ι (left) and the angle θ (right) for GW signals with $f_{\rm gw,1} = 500$ Hz from NSs at $r = 10$ kpc. The left panel of Fig. 2 indicates that the NS mass cannot be estimated at all at $\iota = 0, \pi$ while well estimated at around $\iota = \pi/2$. This is because the amplitudes of the first harmonic mode in both the plus and cross polarizations vanish at $\iota = 0, \pi$ from Eqs. (14a) and (14c). In these cases, since the GW signal has only the second harmonic mode, the effect of the gravitational phase shift is unobservable. The right panel in Fig. 2 indicates that the NS mass cannot be determined for $\theta = 0, \pi/2$. This fact can be interpreted in the same way as the left panel of Fig. 2.



FIG. 1: The cumulative distribution function as a function of relative error of NS mass $\Delta M_{\rm NS}/M_{\rm NS}$ for two different GW frequencies (a) 300 Hz and (b) 500 Hz. The solid line, the dashed line, and the dashed-dotted line corresponds to NSs at the distances $r = 1$ kpc, 10 kpc, and 50 kpc respectively. For NSs at the distance of $r = 50$ kpc that emit GW at the first harmonic GW mode frequency of $f_{\rm gw,1} = 300$ Hz, measurement accuracy becomes more than 100%.



FIG. 2: The scatter-plot of relative error of NS mass $\Delta M_{\rm NS}/M_{\rm NS}$ and inclination ι (left), misalignment angle θ (right). The GW signals are assumed to have the GW frequency 500 Hz from NSs at $r = 10$ kpc.

IV. CONCLUSION

In this paper, we have focused on a gravitational phase shift for a GW from an isolated rapidly rotating non-axisymmetric NS. When the GW passes through the gravitational potential created by the NS, the GW phase is stretched due to the gravitational phase shift effect depending on both its rotational frequency and its mass. We have constructed the explicit form of the resulting phase shift based on Ref. [15] and obtained Eq. (12b). We have shown that if the GW from the NS has different frequency modes such as the first and the second harmonic modes as proposed in [16, 17], the mass of the isolated NS can be, in principle, determined by making use of the phase difference of the different modes. We have given an example of an isolated rapidly rotating non-axisymmetric NS to estimate the measurement accuracy of the NS mass using Fisher analysis in a targeted search. Performing Monte Calro simulations in 3-year ET observation, we have obtained a cumulative distribution function of a relative error of the NS mass $\Delta M_{\rm NS}/M_{\rm NS}$ for different GW frequencies 300 Hz and 500 Hz and different distances of $r = 1, 10$, and 50 kpc as shown in Fig. 1. The range of the NS mass can be typically delimited with an accuracy of $\Delta M_{\rm NS}/M_{\rm NS} = \mathcal{O}(0.1)$ with third generation detectors for such a GW signal that can be detected by a second generation GW detector with SNR $= \mathcal{O}(10)$.

Acknowledgments

We thank Masatake Ohashi for helpful comments. This work is supported by Grant-in-Aid for Japan Society for the Promotion of Science, the JSPS Fellows Grant No. 26.8636 (K. E.), the JSPS Grant-in-Aid for Young Scientists Grant No. 25800126 (Y. I.), and the Ministry of Education, Culture, Sports, Science and Technology (MEXT) Grant-in-Aid for Scientific Research on Innovative Areas "New Developments in Astrophysics Through Multi-Messenger Observations of Gravitational Wave Sources" (Grant Number 24103005) (Y. I.).

Appendix A: GW signals

The GW waveforms from an isolated rapidly rotating non-axisymmetric NS are presented in Sec. III A. As can be seen in Eq. (13a) and (13b), the GW signals consist of two components with frequency $f_{\mathrm{gw},1}$ and $2f_{\mathrm{gw},1}$. It is convenient for our purpose to decompose the GW signals as follows [28].

$$h\left(t\right)=\sum_{j=1}^{2}\sum_{k=1}^{4}A_{jk}h_{jk}\left(t\right). \tag{A1}$$

The amplitudes A_{jk} are defined as

$$A_{11}=h_0\sin2\theta\left[\frac{1}{8}\sin2\iota\cos2\psi\cos\phi_0-\frac{1}{4}\sin\iota\sin2\psi\sin\phi_0\right], \tag{A2a}$$

$$A_{12}=h_0\sin2\theta\left[\frac{1}{4}\sin\iota\cos2\psi\sin\phi_0+\frac{1}{8}\sin2\iota\sin2\psi\cos\phi_0\right], \tag{A2b}$$

$$A_{13}=h_0\sin2\theta\left[-\frac{1}{8}\sin2\iota\cos2\psi\sin\phi_0-\frac{1}{4}\sin\iota\sin2\psi\cos\phi_0\right], \tag{A2c}$$

$$A_{14}=h_0\sin2\theta\left[\frac{1}{4}\sin\iota\cos2\psi\cos\phi_0-\frac{1}{8}\sin2\iota\sin2\psi\sin\phi_0\right], \tag{A2d}$$

$$A_{21}=h_0\sin^2\theta\left[\frac{1}{2}\left(1+\cos^2\iota\right)\cos2\psi\cos\left(2\phi_0+\phi_s\right)-\cos\iota\sin2\psi\sin\left(2\phi_0+\phi_s\right)\right], \tag{A2e}$$

$$A_{22}=h_0\sin^2\theta\left[\frac{1}{2}\left(1+\cos^2\iota\right)\sin2\psi\cos\left(2\phi_0+\phi_s\right)+\cos\iota\cos2\psi\sin\left(2\phi_0+\phi_s\right)\right], \tag{A2f}$$

$$A_{23}=h_0\sin^2\theta\left[-\frac{1}{2}\left(1+\cos^2\iota\right)\cos2\psi\sin\left(2\phi_0+\phi_s\right)-\cos\iota\sin2\psi\cos\left(2\phi_0+\phi_s\right)\right], \tag{A2g}$$

$$A_{24}=h_0\sin^2\theta\left[-\frac{1}{2}\left(1+\cos^2\iota\right)\sin2\psi\sin\left(2\phi_0+\phi_s\right)+\cos\iota\cos2\psi\cos\left(2\phi_0+\phi_s\right)\right] \tag{A2h}$$

where h_0, ι, θ, ψ and ϕ_0 are the overall amplitude, the inclination, the polarization phase, and the initial phase, respectively. The phase ϕ_s expresses the gravitational phase shift due to the gravitational potential of the NS and corresponds to Eq. (15). It is described by

$$\phi_s=8\pi M_{\mathrm{NS}}f_{\mathrm{gw},1}\ln2. \tag{A3}$$

The time-dependent parts $h_{jk}\left(t\right)$ in Eq.(A1) characterize the shape of the GW signal and are defined as

$$h_{j1}\left(t\right)=a\left(t\right)\cos\left[j\Phi\left(t\right)\right], \tag{A4a}$$

$$h_{j2}\left(t\right)=b\left(t\right)\cos\left[j\Phi\left(t\right)\right], \tag{A4b}$$

$$h_{j3}\left(t\right)=a\left(t\right)\sin\left[j\Phi\left(t\right)\right], \tag{A4c}$$

$$h_{j4}\left(t\right)=b\left(t\right)\sin\left[j\Phi\left(t\right)\right] \tag{A4d}$$

where the function $a\left(t\right)$ and $b\left(t\right)$ are modulation amplitudes given in [28]. The GW phase after the barycentric corrections is expressed by

$$\Phi\left(t\right)=2\pi f_{\mathrm{gw},1}\left(t+\boldsymbol{n}\cdot\boldsymbol{r}\left(t\right)\right) \tag{A5}$$

where $\boldsymbol{n}$ denotes a unit vector pointing toward the NS from the solar system barycenter (SSB), and $\boldsymbol{r}\left(t\right)$ denotes the separation vector pointing from the detector to the SSB. Note that we neglected the spin-down effect in the GW frequency for simplicity. The decomposition of the GW signals in Eq.(A1) separates known parameters and unknown parameters in targeted searches.

Appendix B: Fisher analysis

In this appendix, we give a brief review of the Fisher analysis [30, 31]. The detector output $s\,(t)$ can be expressed by linear sum of the GW signal $h\,(t)$ and the detector noise $n\,(t)$, $s\,(t) = h\,(t) + n\,(t)$. When the noise is stationary and obeys the Gaussian distribution, measurement errors of the waveform parameters $\boldsymbol{\lambda}$ are found by

$$\left(\Delta\lambda^j\right)_{\rm rms} \equiv \sqrt{\left\langle \left(\Delta\lambda^j\right)^2\right\rangle} = \sqrt{\left(\Gamma^{-1}\right)_{jj}} \tag{B1}$$

for large signal-to-noise ratio where $\left(\Delta\lambda^j\right)_{\rm rms}$ is the root-mean square of the waveform parameters $\Delta\lambda^j$. The matrix Γ_{jk} is called the Fisher information matrix and is defined by

$$\Gamma_{jk} = \left(\partial_j h \middle| \partial_k h\right) \tag{B2}$$

where $(\cdot|\cdot)$ denotes the noise-weighted inner product, and $\partial_j h$ denotes the derivative of the GW signal with respect to the waveform parameter λ_j. For instance, the inner product between two time functions $x\,(t)$ and $y\,(t)$ is expressed by

$$(x|y) = 4{\rm Re}\int_{-\infty}^{\infty} \frac{\tilde{x}\,(f)\,\tilde{y}^*\,(f)}{S_n\,(f)}df. \tag{B3}$$

where $S_n(f)$ is the one-sided powerspectral density of the detector. In this paper, we have adopted $S_n(f)$ of ET given in the Table 1 of [32].

When the frequency of the signal is nearly constant, it is convenient to introduce a new inner product

$$(x||y) \equiv \frac{2}{T_{\rm obs}}\int_{-T_{\rm obs}/2}^{T_{\rm obs}/2} x\,(t)\,y\,(t)\,dt \tag{B4}$$

where $T_{\rm obs}$ denotes the observation time [28]. Since the time scales of change in modulation amplitudes $a\,(t)$ and $b\,(t)$ are of the order of the time scale of the Earth's rotation, the GW phase $\Phi\,(t)$ oscillates much more rapidly than modulation amplitudes. Then, the inner product between the amplitudes h_{jk} can be well-approximated as

$$(h_{j1}||h_{j1}) \simeq (h_{j3}||h_{j3}) \simeq \frac{1}{2}\,(a||a)\,, \tag{B5a}$$

$$(h_{j2}||h_{j2}) \simeq (h_{j4}||h_{j4}) \simeq \frac{1}{2}\,(b||b)\,, \tag{B5b}$$

$$(h_{j1}||h_{j2}) \simeq (h_{j3}||h_{j4}) \simeq \frac{1}{2}\,(a||b)\,. \tag{B5c}$$

Combining Eqs. (B2)-(B5c), we obtain the Fisher matrix in our analysis,

$$\begin{aligned}\Gamma_{jk} = \sum_{\ell=1,2}\frac{T_{\rm obs}}{2S_n\left(\ell f_{{\rm gw},1}\right)}\Big[&\left(\partial_j A_{\ell 1}\partial_k A_{\ell 1} + \partial_j A_{\ell 3}\partial_k A_{\ell 3}\right)(a||a) + \left(\partial_j A_{\ell 2}\partial_k A_{\ell 2} + \partial_j A_{\ell 4}\partial_k A_{\ell 4}\right)(b||b)\\ &+ \left(\partial_j A_{\ell 1}\partial_k A_{\ell 2} + \partial_j A_{\ell 2}\partial_k A_{\ell 1} + \partial_j A_{\ell 3}\partial_k A_{\ell 4} + \partial_j A_{\ell 4}\partial_k A_{\ell 3}\right)(a||b)\Big].\end{aligned} \tag{B6}$$

---

[1] F. Weber, ed., *Pulsars as astrophysical laboratories for nuclear and particle physics* (1999).

[2] J. M. Lattimer, Annual Review of Nuclear and Particle Science **62**, 485 (2012).

[3] F. M. Walter, S. J. Wolk, and R. Neuhäuser, Nature (London) **379**, 233 (1996).

[4] J. A. Pons, F. M. Walter, J. M. Lattimer, M. Prakash, R. Neuhäuser, and P. An, Astrophys. J. **564**, 981 (2002).

[5] J. M. Lattimer, Astrophys. Space Sci. **336**, 67 (2011).

[6] J. Abadie, B. P. Abbott, R. Abbott, M. Abernathy, T. Accadia, F. Acernese, C. Adams, R. Adhikari, P. Ajith, B. Allen, et al., Classical and Quantum Gravity **27**, 173001 (2010).

[7] Y. Aso et al. (KAGRA Collaboration), Phys.Rev. **D88**, 043007 (2013).

[8] G. M. Harry and LIGO Scientific Collaboration, Classical and Quantum Gravity **27**, 084006 (2010).

[9] F. Acernese, M. Agathos, K. Agatsuma, D. Aisa, N. Allemandou, A. Allocca, J. Amarni, P. Astone, G. Balestri, G. Ballardin, et al., Classical and Quantum Gravity **32**, 024001 (2015).

[10] M. Punturo, M. Abernathy, F. Acernese, B. Allen, N. Andersson, K. Arun, F. Barone, B. Barr, M. Barsuglia, M. Beker, et al., Classical and Quantum Gravity **27**, 194002 (2010).

[11] S. Hild, M. Abernathy, F. Acernese, P. Amaro-Seoane, N. Andersson, K. Arun, F. Barone, B. Barr, M. Barsuglia, M. Beker, et al., Classical and Quantum Gravity **28**, 094013 (2011).

[12] N. Andersson, V. Ferrari, D. I. Jones, K. D. Kokkotas, B. Krishnan, J. S. Read, L. Rezzolla, and B. Zink, General Relativity and Gravitation **43**, 409 (2011).

[13] A. Messiah, *Quantum mechanics* (Dover Publications, 2014).

[14] L. Blanchet and T. Damour, Phys.Rev. **D46**, 4304 (1992).

[15] H. Asada and T. Futamase, Phys.Rev. **D56**, 6062 (1997).

[16] M. Zimmermann and E. Szedenits, Phys.Rev. **D20**, 351 (1979).

[17] D. I. Jones, Mon. Not. R. Astron. Soc. **402**, 2503 (2010).

[18] K. Makishima, T. Enoto, J. S. Hiraga, T. Nakano, K. Nakazawa, S. Sakurai, M. Sasano, and H. Murakami, Phys. Rev. Lett. **112**, 171102 (2014).

[19] K. Makishima, T. Enoto, H. Murakami, Y. Furuta, T. Nakano, M. Sasano, and K. Nakazawa, ArXiv e-prints (2015), 1501.07684.

[20] D. I. Jones and N. Andersson, Mon. Not. R. Astron. Soc. **324**, 811 (2001).

[21] D. I. Jones and N. Andersson, Mon. Not. R. Astron. Soc. **331**, 203 (2002).

[22] D. I. Jones, Mon. Not. R. Astron. Soc. **420**, 2325 (2012).

[23] M. Bejger and A. Królak, Classical and Quantum Gravity **31**, 105011 (2014).

[24] L. D. Landau and M. E. Lifshits, *The classical theory of fields* (Butterworth-Heinemann, 1975).

[25] C. Cutler, Phys. Rev. D **66**, 084025 (2002).

[26] M. Maggiore, *Gravitational Waves: Volume 1: Theory and Experiments* (Oxford University Press, Oxford, 2007).

[27] D. I. Jones, ArXiv e-prints (2015), 1501.05832.

[28] P. Jaranowski, A. Królak, and B. F. Schutz, Phys.Rev. **D58**, 063001 (1998).

[29] C. J. Horowitz and K. Kadau, Physical Review Letters **102**, 191102 (2009).

[30] L. S. Finn, Phys.Rev. **D46**, 5236 (1992).

[31] C. Cutler and E. E. Flanagan, Phys.Rev. **D49**, 2658 (1994).

[32] B. S. Sathyaprakash and B. F. Schutz, Living Reviews in Relativity **12**, 2 (2009).

[33] There is no Coulomb phase shift when the GW leaves the pulsar at $r = 0$.